

Week 16: April 15-21, 2012

ILLINOIS DEPARTMENT OF PUBLIC HEALTH



Illinois Influenza Surveillance Report

Week 16: Week Ending Saturday, April 21, 2012

Division of Infectious Diseases Immunizations Section

4/27/2012

Contents

Summary.....	3
CDC Flu View.....	4
ILI Net Provider Surveillance	5
ILI Visits by Age Group.....	6
Great Lakes Naval Recruit Influenza Surveillance	7
Influenza Intensive Care Unit Admissions and Deaths.....	7
Influenza Related ICU Admissions by Age Group.....	8
Laboratory Surveillance	9
Viral Resistance	10
Influenza Outbreaks Reported in Long-Term Facilities (LTC) and Nursing Homes (NH)	10
IDPH, Immunization Section Regional Map	11
Weekly Viral Subtype	12
Resources.....	13

Summary

- During CDC week 16, the proportion of outpatient visits for influenza-like illness (ILI)¹ was 1.9% compared with 2.0% for week 15.
- Based on CDC criteria, influenza activity is classified as **sporadic** (see CDC FLU View Section) for week 16. This represents no change in activity from week 15.
- Febrile Respiratory Illness (FRI) surveillance² at Naval Recruit Training Command, Great Lakes was **at or below expected value** for week ending April 21, 2012.
- During week 16, none of the two specimens tested by Illinois Department of Public Health Laboratory was positive for influenza.
- No influenza-associated Intensive Care Unit (ICU) admission³ were reported for week 16.
- No influenza-associated pediatric deaths were reported for week 16.
- During week 16, three influenza outbreaks were reported in long-term care facilities within the Peoria (2) and West Chicago (7) regions (see IDPH, Immunization Section Regional Map).

¹ ILI "Influenza like Illness" is defined as fever $\geq 100^{\circ}\text{F}$ and cough and/or sore throat.


² FRI surveillance is ongoing at 8 U.S. military basic training centers, representing all service branches. FRI Rate Status is classified into one of 3 categories:

1. At or below expected value (expected value shown as dashed line)
2. Moderately elevated
3. Substantially elevated

³ For the purpose of diagnosis, influenza can be diagnosed by using the following test: reverse transcription polymerase chain reaction [RT-PCR], viral culture, Immunofluorescence [Direct Fluorescent Antibody (DFA) or Indirect Fluorescent Antibody (IFA) Staining], Enzyme Immuno Assay (EIA) or any rapid diagnostic test. Sensitivities of rapid diagnostic tests are approximately 50-70% when compared with viral culture or reverse transcription polymerase chain reaction (RT-PCR), and specificities of rapid diagnostic tests for influenza are approximately 90-95%. False-positive (and true-negative) results are more likely to occur when disease prevalence in the community is low, which is generally at the beginning and end of the influenza seasons. False-negative (and true-positive) results are more likely to occur when disease prevalence is high in the community, which is typically at the height of the influenza season.


ILI Net Provider Surveillance

Influenza Like Illness Outpatient Surveillance 2010-2012



ILI Visits by Age Group

2010 -12 INFLUENZA SEASON PROPORTION OF ILI OFFICE VISITS BY AGE GROUP



Great Lakes Naval Recruit Influenza Surveillance

Febrile Respiratory Illness (FRI) surveillance⁴ at Naval Recruit Training Command, Great Lakes was **at or below expected value** for week ending April 21, 2012. For more information visit <http://www.med.navy.mil/sites/nhrc/geis/Pages/default.aspx>

Influenza Intensive Care Unit Admissions and Deaths


There were no influenza related ICU admissions or deaths reported for week ending April 21, 2012.

Year	Week No	Admissions No	Deaths
2012	3	1	0
2012	4	1	0
2012	5	0	0
2012	6	1	0
2012	7	3	0
2012	8	4	0
2012	9	2	0
2012	10	14	0
2012	11	9	0
2012	12	4	0
2012	13	8	0
2012	14	5	0
2012	15	4	0
2012	16	0	0

⁴ FRI surveillance is ongoing at 8 U.S. military basic training centers, representing all service branches. FRI Rate Status is classified into one of 3 categories:

4. At or below expected value (expected value shown as dashed line)
5. Moderately elevated
6. Substantially elevated

Influenza Related ICU Admissions by Age Group



Laboratory Surveillance

During week 16, two specimens were tested by Illinois WHO/NREVSS^[1] collaborating laboratories in Illinois. This represents an decrease in testing compared with week 15. None of the two specimens tested by Illinois Department of Public Health Laboratory was positive for influenza.

Year	Week	A (H1)	2009(A)H1N1	A (H3)	A(Unable to subtype)	A(Sub typing not performed)	B	Total # Tested	% Positive
2012	1	0	0	3	0	0	1	4	100%
2012	2	0	0	3	0	1	0	7	57%
2012	3	0	1	3	0	1	1	10	60%
2012	4	0	0	2	0	0	0	4	50%
2012	5	0	0	0	0	0	0	4	0
2012	6	0	3	1	0	0	1	9	56%
2012	7	0	0	3	0	0	0	4	75%
2012	8	0	6	13	0	0	4	30	77%
2012	9	0	12	12	0	0	1	30	83%
2012	10	0	4	8	0	0	0	30	40%
2012	11	0	11	24	0	0	0	46	76%
2012	12	0	2	3	0	0	0	9	56%
2012	13	0	0	4	0	0	0	11	36%
2012	14	0	1	0	0	0	0	2	50%
2012	15	0	0	1	0	0	0	4	25%
2012	16	0	0	0	0	0	0	2	0

^[1] WHO/NREVSS Collaborating Laboratories WHO/NREVSS Collaborating Laboratories: For viral surveillance - About 80 U.S. World Health Organization (WHO) Collaborating Laboratories and 70 National Respiratory and Enteric Virus Surveillance System (NREVSS), located throughout the United States report daily or weekly the results of their testing.

Viral Resistance

Neuraminidase Inhibitor Resistance Testing Results on Samples Collected Since October 1, 2011


	Oseltamivir		Zanamivir	
	Virus Samples tested (n)	Resistant Viruses, Number (%)	Virus Samples tested (n)	Resistant Viruses, Number (%)
Influenza A (H3N2)	1,072	0 (0.0)	1,072	0 (0.0)
Influenza B	217	0 (0.0)	217	0 (0.0)
Influenza A (2009 H1N1)	682	15 (2.2)	393	0 (0.0)

Rare sporadic cases of oseltamivir resistant 2009 H1N1 and A (H3N2) have been detected worldwide. Antiviral treatment with oseltamivir or zanamivir is recommended as early as possible for patients with confirmed or suspected influenza who have severe, complicated, or progressive illness; who require hospitalization; or who are at greater risk for influenza-related complications. Additional information on recommendations for treatment and chemoprophylaxis of influenza virus infection with antiviral agents is available at <http://www.cdc.gov/flu/antivirals/index.htm>.

Influenza Outbreaks Reported in Long-Term Facilities (LTC) and Nursing Homes (NH)

Three influenza outbreaks were reported in long-term care facilities within the Peoria (2) and West Chicago (7) regions (see IDPH, Immunization Section Regional Map); bringing the total number of outbreaks reported during 2011-12 Influenza season to 36.

Region	Number of outbreaks No. (%)
Rockford (1)	6(17)
Peoria (2),	5(14)
Edwardsville (4),	6(17)
Marion (5),	3(8)
Champaign (6),	0(0)
West Chicago (7)	8(22)
Chicago/Cook (8)	8(22)
Total	36(100)




IDPH, Immunization Section Regional Map



Weekly Viral Subtype

Influenza Isolates from Illinois Reported by WHO/NREVSS Collaborating Laboratories 2010-2012



Resources

- Centers for Disease Control and Prevention Influenza Website: <http://www.cdc.gov/flu/>
- Immunization Action Coalition Website: <http://immunize.org/>
- IDPH Website: <http://www.idph.state.il.us/flu/surveillance.htm>
- ACL Clinical Laboratory Respiratory Panel: <http://www.acllaboratories.com/>
- St Louis Children's Hospital Clinical Laboratory Respiratory Panel:
<http://www.stlouischildrens.org/content/ClinicalLaboratories.htm>